







## Asthma Treatment Plan – Adult

			www.pacnj.org	
Name:		Effective Date:		
Doctor:		Phone:		
Follow-up A	Appointment Date:	Time:		
HEALTHY	(Green Zone)	Take daily control medicine more effective with a "space		Triggers Check all items
And/or Peak	You have all of these:  • Breathing is good  • No cough or wheeze  • Sleep through the night  • Can work, exercise, and play	☐ Flovent® Diskus® ☐ 50 ☐ 100 ☐ 250 ☐ Pulmicort Flexhaler® ☐ 90, ☐ 180 ☐		o Pets - animal
	(Yellow Zone)	sthma, take this medicine	e your mouth after taking inhaled medicineminutes before exercise. e(s) and ADD quick-relief medicine(s).	cleaning
9-3-5	You have <u>any</u> of these:  • Cough	MEDICINE	HOW MUCH to take and HOW OFTEN to take it	inside or outside
If quick-relief m 15-20 minutes 2 times and syn doctor or go to	Mild wheeze     Tight chest     Coughing at night     Other: nedicine does not help within or has been used more than mptoms persist, call your the emergency room.  If flow from to	□ Duoneb® Duoneb® 0.31, □ 0.63, □ Combivent Respimat® Increase the dose of, or add: □ Other  • If quick-relief medicine is	2 puffs every 4 hours as needed1 unit nebulized every 4 hours as needed1 unit nebulized every 4 hours as needed ] 1.25 mg _1 unit nebulized every 4 hours as needed	□ Weather ○ Sudden temperature change ○ Extreme weathe - hot and cold ○ Ozone alert days □ Foods: ○ ○
EMERGE	NCY (Red Zone)		es NOW and CALL 911. eatening illness. Do not wait!	O   O
And/or Peak	Quick-relief medicine did no help within 15-20 minutes Breathing is hard or fast Nose opens wide Ribs shows Trouble walking and talking Lips blue Fingernails blue Other:	☐ Combivent® ☐ Maxair® ☐ Xopenex® ☐ Ventolin® ☐ Pro-Air® ☐ Proventil® ☐  OW ☐ Albuterol ☐ 1.25, ☐ 2.5 mg ☐ ☐ Duoneb® ☐	HOW MUCH to take and HOW OFTEN to take it  2 puffs every 20 minutes 2 puffs every 20 minutes 1 unit nebulized every 20 minutes	This asthma treatmer plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.

Disclaimers: The use of this Website/PACNJ Asthma Treatment Plan and its content is at your own risk. The content is provided on an "as is" basis. The American Lung Association of the Mid-Allantic (ALAM-A), the Pediatric/Adult Asthma Coalition of New Jersey and all affiliates disclaim all warranties, express or implied, statutory or otherwise, including but not limited to the implied warranties or merchantability, non-infringement of third parties rights, and filness for a particular purpose. ALAM-A makes no representations or warranties about the accuracy, reliability, completeness, currency, or immeliness of the content. ALAM-A makes no warranty, representation or guaranty that the information will be uninterrupted or error free or that any detects can be corrected. In no event shall, ALAM As he label to any damages, fonding, without limitation, incidental and its affiliates are not liable for any claim, whatsoever, caused by your use or misuse of the Asthma Treatment Plan, nor of this website.

The Pediatric/Adult Ashma Coalition of New Jersey is sponsored by the American Lung Association in New Jersey. This publication was supported by a grant from the New Jersey Department of Health and Senior Services or the U.S. Centers for Disease Control and Prevention under Cooperative Agreement 5U59EH000491-03. Its content are solely the responsibility of the authors and do not necessarily represent the official views of the New Jersey Department of Health and Senior Services or the U.S. Centers for Disease Control and Prevention Although this document has been funded wholly or in part by the United States Environmental Protection Agency under Agreement XA97250908-4 to the American Lung Association in New Jersey, it has not gone through the Agency's publications review process and therefore, may not necessarily reflect the views of the Agency and no official endorsement should be inferred. Information in this publication is not intended to diagnose health problems or take the place of medical advice. For asthma or any medical condition, seek medical advice from your child's or your rhealth care professional.