

2014 UPDATES TO COID GUIDANCE FURTHER RESTRICTS ACCESS TO PROPHYLAXIS

	2014 Guidance	2012 Guidance	Approved Label
≤28 WGA	No Change	No Change	No Change
29-31 WGA	Not recommended unless has other qualifying conditions	Have to be ≤6 months at start of RSV season	Have to be ≤6 months at start of RSV season
32-34 WGA	Not recommended unless has other qualifying conditions	GA defined as: <ul style="list-style-type: none"> • 32 weeks, 0 days through 34 weeks, 6 days, and ≤90 days • AND daycare attendance OR siblings <5 years of age as a risk factor* 	Have to be ≤6 months at start of RSV season
35 WGA	Not recommended unless has other qualifying conditions	Not recommended unless has other qualifying conditions	Have to be ≤6 months at start of RSV season
BPD/CLDP†	<ul style="list-style-type: none"> • CLDP defined as <32 wGA and requiring >21% oxygen for at least the first 28 days after birth; <12 months at RSV season start, medications not required • 12-24 months at RSV season start if still on medications 	≤24 months at RSV season start if requiring oxygen, medications in prior 6 months	≤24 months at start of RSV season
HS-CHD‡	<ul style="list-style-type: none"> • 12-24 months at RSV season start no longer recommended 	≤24 months at RSV season start	≤24 months at RSV season start

Please see Important Safety Information on slides 6 and 7.

*Multiples do not qualify as siblings.

†BPD/CLDP = bronchopulmonary dysplasia/chronic lung disease of prematurity.

‡HS-CHD = hemodynamically-significant congenital heart disease.